

WITCHES BROOM ON PIGEON PEA INDUCED BY MEALY BUG INFESTATIONS.

In January, 1961, 35 out of 56 pigeon pea plants (*Cajanus cajan* Millsp.) in a small plot being grown in the experimental food crop garden at the Lowlands Agricultural Experiment Station, Keravat, New Britain, showed marked witches broom symptoms (Plate II) of the lateral shoots. Each shoot was heavily infested with mealy bugs [*Planococcus citri* (Risso)]† in all stages of development. The plants were thoroughly sprayed with malathion plus wetting agents which eliminated the mealy bugs. Six weeks after spraying, all the plants appeared quite healthy. Buds from abnormal shoots were grafted onto healthy pigeon pea seedlings and of the thirty plants grafted, none developed witches broom shoots.



Plate II.—Witches Brooming of lateral shoots on *Cajanus cajan*.

(Induced by infestations of mealy bugs.)

In the glass house, 20 pigeon pea seedlings were raised. Mealy bugs were collected from stock colonies in the glass house and released onto the seedlings when 10 weeks old. The terminal shoots were removed to induce the development of laterals. After 12 weeks, the lateral shoots had developed the typical witches broom symptoms as observed in the field. Spraying was carried out as in the field trial and all the test plants recovered.

Thus it has been concluded that the witches brooming on *Cajanus cajan* observed at Keravat is induced by heavy infestations of mealy bug [*Planococcus citri* (Risso)].

R. J. VAN VELSON.

(Manuscript received 5th September, 1961.)

† The insect identification was made by the Commonwealth Institute of Entomology.

TWO ANIMAL PARASITES PREVIOUSLY UNRECORDED IN NEW GUINEA.

TWO animal parasites, previously unrecorded in New Guinea were recently identified at Lae. They are *Gongylonema ingluvicola* and *Notoedres cati*.

Gongylonema ingluvicola.

A few specimens of this nematode were found in a mature fowl during a routine post-mortem examination.

G. ingluvicola is found in small burrows in the mucous membrane of the crop of fowls, turkeys and quail. The worm is characterized by a zone of shieldlike markings on the anterior end of the body. Mature male worms measure 1.7-2.0 cm long and mature females 3.2-5.5 cm long.

The life cycle is uncertain but it is thought to involve cockroaches and possibly some species of beetles as intermediate hosts.

The only damage these worms cause are local lesions in the form of burrows in the mucosa of the crop.

It is unlikely that infestation with *G. ingluvicola* alone would cause any ill effects in fowls. Their presence, however, indicates that flock hygiene is faulty and that heavy infestations with other parasites likely. The fowl in which this parasite was identified at Lae was also infected with *Amaebotaenia sphenoides* (a tapeworm), *Capillaria* sp. and *Oxyspirura mansoni* (eye worm), despite treatment of the flock with piperazine.

Notoedres cati.

This parasite was responsible for a severe case of mange in a European owned cat.

N. cati is a small mite which closely resembles *Sarcoptes scabiei* except for the dorsal position of the anus.